

- beta glucan from the cultured fruit body of *Agaricus blazei* [J]. Biol Pharm Bull , 2001 , 24(7) : 820 ~ 828
- 5 Mizuno M , Minato K , Ito H et al. Anti-tumor polysaccharide from the mycelium of liquid-cultured *Agaricus blazei* mil [J]. Biochem Mol Biol Int , 1999 , 47(4) : 707 ~ 714
 - 6 Mizuno M , Morimoto M , Minato K et al. Polysaccharides from *Agaricus blazei* stimulate lymphocyte T-cell subsets in mice [J]. Biosci Biotechnol Biochem , 1998 , 62(3) : 434 ~ 437
 - 7 Nskajima A , Ishida T , Koga M et al. Effect of hot water extract from *Agaricus blazei* Murill antibody-producing cells in mice [J]. International Immunopharmacol 2002 , 2(8) : 1205 ~ 1211
 - 8 Martins de Oliveira J , Jordao B Q , Ribeiro L R et al. Anti-genotoxic effect of aqueous extracts of sun mushroom (*Agaricus blazei* Murill lineage 99/26) in mammalian cells in vitro [J]. Food and Chemical Toxicology 2002 , 40(12) : 1775 ~ 1780
 - 9 Barbisan F L , Miyamoto M , Scolastici C et al. Influence of aqueous extract of *Agaricus blazei* on rat liver-toxicity induced by different dosed of diethylnitrosamine [J]. Journal of Ethnopharmacology , 2002 , 83(1 - 2) : 25 ~ 32
 - 10 杨淑云 , 刘朝贵 , 朱 凯等 . 富硒姬松茸液体培养条件的研究 [J]. 微生物学通报 , 2003 , 30(1) : 9 ~ 13
 - 11 王六生 , 谷文英 . 姬松茸深层发酵培养基的优化 [J]. 无锡轻工大学学报 , 2002 , 21(4) : 389 ~ 392
 - 12 邹 祥 , 张 梁 , 章克昌 . 姬松茸液体培养工艺条件的研究 [J]. 药物生物技术 , 2002 , 9(6) : 339 ~ 343

Influence of the Ages of Inoculums in Submerged Fermentation Process

Dong Xiuping Sun Yumei Li Yanjie Zhu Beiwei

(College of Bio & Food Technology , Dalian Institute of Light Industry , Dalian , 116034)

ABSTRACT Insaccharides , the extra cellular polysaccharides , the extra cellular cellulase , the changes of the reduced sugar and pH in submerged fermentation process were studied. In this paper the effects of inoculums ages on the growth of mycelium , the cellular polysaccharides production and extra cellular polysaccharides were studied. It is seen that inoculums of cultivation for 4 days were higher than those with inoculums of cultivation for 6 days and 8 days. The output of exo-1 β -D-glucanase(enzyme C₁) and endo-1 β -D-glucanase(enzyme C_x) for 8 days were higher than those with inoculums of cultivation for 4 days and 6 days.

Key words *Aaricus blazei* Murril , the ages of inoculums , liquid submerged fermentation

欧美包装市场出现全新的食品专用包装纸

最近 欧美市场又推出了全新的 PLMEX 食品专用包装纸 , 它是 100% 纯纸浆 , 不含荧光剂和危害人体的化学物质 , 具有防水、防油、抗粘、耐热(可耐 200 ~ 250℃ 高温) 的特点。使用方便、清理简单、安全卫生 , 符合美国 FDA 和德国 BGA 食品卫生标准。这种食品专用纸使用后经清水清洗可回收 , 并可以反复使用 50 次。这种食品纸有成卷的 , 也有压制成各种纸杯形状的 , 不管是用于蒸制、烘烤 , 还是微波加热 , 都不会变形和褪色。